# Effect of sowing date on growth, development and yield of gero millet

P. N. EGHAREVBA, S. M. ABED and A. A. OKOLO Institute for Agricultural Research Ahmadu Bello University

### **Abstract**

Zaria.

The objective of this study was to determine the most appropriate time for sowing millet (Pennisetum typhoides) in Samaru. Ex-Bornu millet, an improved local variety, was used and was sown at the currently recommended spacing of 0.28 x 0.75m. Sowing was spread over fourteen weeks (April to July) at biweekly intervals.

Seedling emergence and germination were dependent on the amount of moisture available. There was a curvilinear relationship between time of sowing and 50% heading, indicating a delay in heading when sowing is done too early or too late. Except for the first two sowings, time to 50% heading was positively correlated with accumulated heat units (r =  $0.81**; r_2 = 66%).$ 

Inflection point on the quadratic curve, which was the best fit for grain yield on planting date, corresponded to the May 24th sowings. Every fortnight delay in sowing after this date resulted in a yield reduction of 294 kg ha-1. Grain yield was positively correlated with established number of plants per unit land area (r =  $0.92^{48}$ ;  $r^2 = 85\%$ ).

Millet should be sown after the first 5 - 8 rains of the season.

#### Introduction

Millet (Pennisetum typhoides S. and H), in spite of its importance as the basic daily diet of many people in Africa and India, continues to receive very low attention in terms of research. While considerable work has been done on the influence of sowing date on the growth, maturity and yield of maize (Zea mays L.) (Zuber, 1966; Bunting, 1968; Genter and Jones, 1970) and of sorghum Sorghum bicolor L.) (Stickler and Pauli, 1966; Kassam and Andrews, 1975; Millington et al., 1977), very little information is available on the effect of sowing date on the performance of millet. This has led to the habit of many farmers sowing with the first rain of the season. This first rain of the year does not follow any set pattern, it could fall as early as February or as late as May. Thus, this chance sowing with the first rain succeeds in only about 8% of the time ( Egharevba 1979 ). Farmers usually give two reasons for sowing with the first rain. Firstly, they claim that the millet sown gives greater yield and has fewer pest problems. Secondly, the millet matures early enough to provide a source of food when other crops like maize and sorghum are still growing. These not withstanding, chance sowing with the first rain will only continue to give an overall low vield of millet. Only experimentally determined time of

sowing can give an increase in yield. In spaper reports the effects of sowing date on growth, development and yield of Gero millet.

### **Meterials And Methods**

The experiment was conducted during the wet seasons of 1976 to 1978 in Samaru (11° 11′N, 7°; 38′E), Nigeria. Ex-Bornu millet, an improved local variety, was used in the trials which were laid out as randomized complete block designs with four replicates. The currently recommended spacing of 0.28 x 0.75m (equivalent to 50,000 plants per hectare) was used. Each plot consisted of 8 rows spaced at 0.75m apart and each row was 6m long for a total plot size of  $_{36~m}^2$ . The four inner rows were harvested for total yield determination.

Soils of the experimental area have been previously described by Jones (1973) Fertilizer (N and P<sub>2</sub>0<sub>5</sub> equivalent to 75 and 30 kg ha—1 respectively) was applied at planting.

Seven sowing dates, designated d1-d7 (Table 1), were used. Sowing started each year on 12th April, and continued at 2-week intervals for a total of 7 plantings. At each planting, seeds were oversown but thinned to one plant per stand two weeks later. Weeds were controlled by hoeing while plants were sprayed periodically with carbaryl 1 at the rate of 1.12 kg ai/ha to control pests.

Observations and measurements made included: percent emergence, number of plants established, time to 50% heading and plant height. Accumulated heat unit was calculated according to the formula:—
(Minimum temp + Maximum temp) — base temperature (Amold, 1960). Base

temperature for millet was taken as 10oC (Kumar, personal Communication).

### **Results and Discussion**

# Emergence

Time to emergence, percent emergence and rainfall data are shown in Table 1. Time to emergence ranged from 5 to 13 days and was significantly longer in those sowing dates in which rainfall was below 50mm. Sowing dates with few rain days delayed emergence.

Percent emergence was significantly (P / .05) lower for the first two sowing dates in both 1976 and 1978. The first two sowings were not done in 1977 because of lack of rainfall, but percent emergence in d<sub>3</sub> and d<sub>4</sub> were significantly (P<.05) lower than others that year.

The delay in emergence of seedlings at the earlier sowing dates (dr and d2) had to do with inadequate supply of moisture. Water is essential for

<sup>1</sup> Formulated as Vetox 85 (Shell Chemical Co.)

TABLE 1. THE EFFECT OF SOUTH DATE OF MELLET ON TIME TO EMERGENCE, & GERMINATION AND PLANT ESTABLISHMENT

			1976				1977			
SowIng Date	Days to Emerg.	Percent Emerg.	P1. Estab. /m <sup>2</sup>	Total Rainfall (mm)	in. of Pain Jays	Days to Emerg.	Percent Emerg.	PI+, Estab.	Total Rain- fall(mm)	No. of Rain days
12 Apr. (d <sub>1</sub> )	10	64	9	26.4	2	-	-	-	-	-
26 Apr. (d <sub>2</sub> )	8	68	13	41.0	3	-	-	-	-	-
IO May (d <sub>x</sub> )	6	89	20	100.1	6	13	41	11	26.3	3
24 May (d <sub>4</sub> )	5	93	22	169.8	3	11	58	13	31.2	4
7 June (d <sub>5</sub> )	5	90	15	220.4	5	6	84	19	132.3	6
21 June (d <sub>6</sub> )	6	92	8	359.4	7	7	90	12	165.7	7
5 July (d <sub>7</sub>	6	90	4	502.6	6	5	89	6	182.3	. 6
Mean	6.6	83.7	13	202.8	5.4	8.2	72.6	12.2	107.6	5.2
S E	0.55**	3.50**	1.4**	-	-	.65**	2.89**	1.17 <sup>ns</sup>	-	-
LSD (.05)	2.0	11.0	5.0	-	•	2.0	10.0	-	-	

<sup>\*\*</sup> Significant at .01

<sup>\*</sup> Significant at .05

ns Not significant at 0.5

TABLE 1 ..... Contd.

Sowing Date	Days to	1978 Percent	PIt. Estab	. Total	al No. of	Sowing Date	The t	Days to	Percent	7	76, 77 & 78) PI. Estab.	
To Talling	Days to Emerg.		PIt. Estab. per m <sup>2</sup>	~	- 7 0	Parallel Local	E <sub>m</sub> Da		Percen Emerg.	cent rg.	t PI. Estab.	cent Pl. Estab. Total rg. per m 2 Rain- fall (mm)
12 Apr.(d,)	9	68	10	26.9	9 3	12 Apr. (0, *)	9	9.5	0	66	9,5	9,5
26 Apr. (d,)	00	72	13	48.7	7 2	26 Apr. (d,+)	00	8.0		70	70 13.0	
10 May (d2)	5	89	. 19	107.6	6 5	IO May (d <sub>3</sub> )	00	8.0		73	73 16.7	16.7
24 May (da)	5	92	22	172.5	5 6	24 May (da)	7	7.2	4	81	19.0	19.0
7 June(d <sub>5</sub> )	6	90	16	298.6	6 8	7 June(d <sub>5</sub> )	- -	5.7		88	88 16.7	16.7
21 June(d <sub>6</sub> )	5	92	01	392.6	6 4	2 June(d <sub>6</sub> )	OI OI	5.7		91	91 10.0	10.0
5 July(d,)	6	92	5	450.3	3 4	5 July(d7)	UT	5.8		90	90 5.0	
Mean	6.3	85.0	13.6	213.9	9 4.6	G. Mean	6	6.9	~	81.3	31.3 13.0	
S. E.	.48**	2.91**	1.18**	ī		S.E (n=9)	0	0.32**		1.85**	1.85** o.75**	
0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	2.0	9.0	4.0	,		S E(n=6)	0	0.39		2.26	2.26 0.91	

germination and if adequate moisture is not available, germination may either be delayed or fail to occur. This is substantiated by the findings of Zuber (1968) for maize and Evans and Stickler (1961) for sorghum, that partial wetting of seeds delayed germination and that the resultant seedlings were weak, giving poor stands. Similar poor stand establishment was observed for dr and d2 (Table 1) in this study; and was also evident in the later sowings (d6 and d7), despite the fact that there were no significant differences in percent emergence from d3 to d7. Thus, there is a need to distinguish between seedling emergence and stand establishment. While seedling emergence is the appearance above ground of germinated seeds, stand establishment is the ability of the emerged seedling to develop and produce harvestable head. The lower stand establishment for the late sowings could be explained partly by progressive reduction in accumulated heat unit and reduction in available nutrient as a result of leaching.

# Developmental phases

Number of days from sowing to heading, plant height and accumulated heat unit associated with the 14 week spread in dates of sowing are shown in Table 2. In 1976, heading occurred 58 days after planting in the first sowing while it occurred after 62 days in the 7th. The earliest time to heading was 53 days while the latest was 62, giving a total spread of 9 days. Similar spread for 1977 and 1978 were 6 and 9 days respectively, giving a mean spread of 8 days for the three years.

Plant height ranged from 162 cm in 1977 to 243 cm in 1976 for a mean of 209 cm and an average of 12% coefficient of variation over the three years. Accumulated heat units decreased with planting date indicating a general decrease in temperatures as the growing season progressed. There was a decrease of 48% in heat units between the first and the last sowing date in 1976. Corresponding decreases for 1977 and 1978 were 37 and 46% respectively. The overall decrease in heat units between the first and last plantings during the three years was 43%.

Days to 50% heading demonstrated a quadratic relationship with sowing date (Fig. 1). This indicates that date of heading is delayed by either planting too early or too late. Point of inflection in Fig. 1 is d4 (24 May). The regression line from first sowing date (10 April) to the point of inflection shows a negative linear function with correlation coefficient,  $r = -0.91^{**}$  ( $r^2 = 83\%$ ) and a regression equation,  $Y_1 = 60.12 - 1.85X_1$  (Fig. 2). This means that there was a delay of 1.85 days in time of heading for each fortnight of sowing earlier than 24 May. Similarly, the regression line from point of inflection to the last sowing date (5 July) shows a positive linear function with a correlation coefficient,  $r = 0.93^{**}$  ( $r^2 = 84\%$ ) and a regression equation,  $Y_1 = 41.37 + 2.77X_1$ . This again indicates that for each fortnight delay in sowing after 24 May there is a delay of 2.77 days in time to heading. This supports the findings of Bonaparte and Brawn (1976) for maize and Kassam and Andrews (1975) for sorghum that delay in sowing after establishment of rains leads to a delay in heading date.

TABLE 2. THE EFFECT OF SOMING DATE AND ACCUMULATED HEAT UNIT ON TIME TO SOM HEADING AND PLANT HEIGHT IN MILLEY

		976			1761	710
Sowing Date	Days to 50% heading	Pi. ht.	Accum. ht. unit	Days to 50% heading	PI. ht. (cm)	Accum. ht. unit
April 12th (d,)	58	202	1524	lon test	a sile sile sile sind win	TOTAL SECTION
April 26th (d2)	57	215	1392	inc ab	- 0	arti arti dis
Way 10th (dz)	53	222	1158	95	194	1552
May 10th (da	53	243	1040	54	218	1319
June 7th (dg)	55	238	978	53	229	1150
June 21st (dg)	65	961	892	22	183	1080
July 5th (dy)	62	176	788	65	162	984
Nesin nesin	56.7	213.1	.0111	55.8	197.2	1217.0
3:E	1,65*	7.99	73.58**	1.62 <sup>ns</sup>	11.77 <sup>ns</sup>	*65.69
L.S.D. (0.5)	5.0	25	727			277

# Significant at .01 # Significant at .05 ns not Significant at .05

TABLE 3 (Contd.

Constant Date		1978			Sowing Datee	The three years (1976, 77 and 78) Combined	years (	77 ,9761)	and 78	Combined
Di Di	Days to 50% heading	PI. ht.		Accum. heat unit	Clear that a	Days to 50% heading	A	PI. ht. (cm)	he	Accum. heat unit
12 Apr. (d,)	59.	208		1508	12 Apr.(d,+)	58.5		205	:	1516
26 Apr. (d2)	58	218		1400	26 Apr. (d2+)	57.5		216		1396
ID May (dz)	54	225	A Million	1242	IO May (dz)	54.4		214		1317
24 May (da)	53	242		1166	24 May (d <sub>a</sub> )	53.3	0.00	324		1175
June(dg)	53	237		1156	7 June(dg)	53.7		235		1095
21 June(dg)	65	200		962	21 June (dg)	58.3		193		798
SSJuly(d <sub>7</sub> )	29	167		842	5 July(d7)	0.19		168		871
Mean	56.8	213.8		1182,3	G. Mean	56.6		209.2	_	1164.9
S. E.	0.94**	10.49**	E	54.16**	S.E. (n=9)	0.82**		5,77**		38.04**
L.S.D. (.05,	3.0	33		167	S.E. (n=6)	00.1		7.06		46.59

\*\* Significant at .01
\* Significant at .05

+  $d_1$  and  $d_2$  are based on 6 observations  $d_3$  to  $d_7$  are based on 9 observations

ns Not Significant at .05

TABLE 3. CORRELATION CORPECTENT, FMATRIX OF SONING DATE  $(d_3-d_4)$  and Different traits in MILLET

- 12 - 12 - 12 - 12 - 12 - 12 - 12 - 12	Estat Horner	••••									
	70)\$ Heading	•	-								
	Pl. helgh:	;		• • • • • • • • • • • • • • • • • • • •							
	1000 -MT	•		·	* * * * * * * * * * * * * * * * * * * *						
	Gays to emer				130.5.	:					
-, signal	Syreleation			.s.u.,	. S. n . C.	:	• • • • •				
	Rainfall				. \$. <sup>8</sup> . 1. 5.	:: :::::::::::::::::::::::::::::::::::	•	*,*1			
.e6777 .9169	Rainy Days	•		-5-647-	.3. n.s.	·	•	• • • • • • • • • • • • • • • • • • • •	e: 		
Grain Establish 50%, Plant 1988-41 Days to Percent Total Yield ment Heading Helght Helly: Therrence Emer Dainfall	Hoat Unit	c .		• · · · · ·	e.n.c.	S. C	•	- ##ct	: i	:	
Establish 50% Plant 1999 at Days to Percent Total ment weating Height relight Themence From Spinis!!	Cowing Date	*****		; 6.	. 64.	٠	. s. " s.	• ;;	•	<i>.</i>	: :
*+04+6"		Grain Yield	ш	50€. Heading	Plant Helght	1973 - 41	Tays to P.	ercert Te	otal Sisfall		
					,		Ā,	rc.,			

\*\* Significant at .0]

\* Significant at .05

ns Not significant at .05.

TARLE 4. THE EFFECT OF SOMING DATE ON GRAIN VIELD AND 1000 KERNEL WEIGHT OF MILLET

Ċ

	1336		1977		1978		All three Years Combined 1974, 71 & 74	Combined 1976,	HL 8 TF
Sowing Date	Grain Yield kg/ha.	1000 K. ₩↑. (grams)	Grain Yield 1000 K. Grain Vield kg/ha. wt. kg/ha (grams)	100 K. ₩+. (grams)	Grain vield kg/ha	100 K.	Sowing Cate	Grain Vield Rô/ha	100 K. **. (grams)
12 Apr. d,	5.50	9.01	1	,	523	0.01	12 Apr. (d,+)	1. 8,6	19.7
ie Apr. d2	ر. رياه	10.2	ı	,	749	10.2	26 Apr. (d3+)		<u></u>
10 May da	S: -	10.4	\$ <u>}</u> £	œ. •	الغ2ن	12.2	(*F) ARM ():	कंकिं∷  -	
34 May d	( ) a	F. 0	i.	10.2	0:1	·:-	74 May (d.)	1 25.7	, <u>.</u>
7 June de	<u>G</u>	8.01	(::	<u>.</u>	(	· .	( b) Jung .	7	·: `-
1 June of	<del>-</del> -	10.4	4	1:51	•a.		C. June (C.)	e . ?	. : 
ئا بادار ہ	о 4	¥ .5	£23	or c	.;	-	("O) Ainf ;	7	
ued.		#; -	::	<u>:</u>		:	ca e		-
ui 	• 7. · Tr	0.349.9	:	V	•			•	1,210
S. D. 1. C. S. 7	<i>(</i> ,	,	( , <u>e</u>	•		•	S.F. (n=6)		<u>a</u> .

\*\* Significant at .01

+  $\mathbf{d}_1$  and  $\mathbf{d}_2$  are based on 6 observations  $\mathbf{d}_1$  to  $\mathbf{d}_2$  are based on 9 observations.

8

TABLE S. CORRELATION COEFFICIENT, r MATRIX OF SOWING DATE  $(d_{\frac{1}{2}}-d_{\frac{1}{2}})$  AND DIFFERENT TRAITS IN MILLET

	Grain Yield	Establish- ment	50% Heading	Plant Height	1000 K Weight	Days to Emergence	Percent Emergence	Total Rainfail	No. of Rain days	Accum. ht. unii
Sowing Date	10 <sup>n.s</sup>	35 <sup>n.s</sup>	.31 <sup>n.s</sup>	46	.n2 <sup>n.s</sup>	~.55 <sup>**</sup>	.62	.82	.55**	90
Heat Unit	09 <sup>n.s</sup>	.21 <sup>n.s</sup>	28 <sup>n.s</sup>	.28 <sup>n.s</sup>	.16 <sup>n.s</sup>	. 76	83	88	62	
No. Rainy days	.53**		37 <sup>n.s</sup>		.23 <sup>n.s</sup>	64	.69	.41	**	
T-Rainfail	10 <sup>n.s</sup>	41	.48	37n.s	.02 <sup>n.s.</sup>		.67			
<b>≰</b> Germination	.44	.16 <sup>n.s</sup>	.01 <sup>n.s</sup>	.05 <sup>n.s</sup>	.33 <sup>n.s</sup>	97				
Days to emerg.	48	24 <sup>n.5</sup>		-12 <sup>n.s</sup>	34 <sup>n.s</sup>					
1000 KWT	.60**	.52**	37*	.59**						
Pl. Height	.84**	.88**	84**							
50 <b>% Headin</b> g	82**	92**								
Establishment	.92**									

<sup>\*\*</sup> Significant at .01

<sup>\*</sup> Significant at .05

ns Not significant at .05




Fig. 1: The quadratic effect of sowing date on time to 50%, heading in millet




Fig. 2: The linear effects of sowing date on time to 50% heading in millet

These studies however did not consider the effect of sowing too early. The delay in heading date due to either sowing too early or too late can be explained partly by drought effect and partly by the decrease in accumulated heat units over the season. Millet, even though drought tolerant, requires certain critical minimum amount of moisture for optimum development. From the rainfall data (Table 1), it would seem that minimum rainfall requirement was not met at the April sowing dates and this could have slowed down the developmental processes. With the exclusion of the two April sowing dates, there was a highly significant negative correlation (r =-0.65;  $r^2 = 42\%$ ) between total heat units accumulated and time to heading (Table 3). Heading time increased as heat unit decreased.

#### Grain Yield

The grain-yield data are shown in Table 4. The lower yield recorded in 1977 is evident from Fig. 3 and this certainly had to do with the rainfall pattern Rainfall was lower and did not become constant until about the first week in June (Table 1). Consequently, peak production was recorded for the sowing made in the first week of June (d5). Subsequent sowings led to a significant reduction in yield.

In 1976 and 1978, best yields were obtained from sowings made in May which coincided with the time when rainfall was regarded as being constant. This is so because while there was a total of 5 rain days up till 26 April of each year for a total of less than 50 mm, there were 15 (total of 128,8 mm) and 11 (total of 122.8 mm) rain days, respectively between 26 April and 24 May in 1976 and 1978. Averaged over the three years, sowings made on 24 May

gave the highest yield.

Sowings made before and long after the rains were steady resulted in significant reduction in yield (Table 4). Grain yield, averaged across years, demonstrated a significant quadratic relationship with time of sowing (Fig. 4). This means that grain yield increased with each successive extension in date of sowing, got to a peak (inflection point on Fig. 4) and then declined. The first half of the curve is characterized by a positive linear function with a correlation coefficient, r = 72 (r2 = 52%) and a regression equation, Yi = 316.0 + 269.5Xi (P∠05); indicating that yield increased by 269.5 kg ha-1 for each fortnight delay in planting. On the contrary, the second half of the curve is characterized by a negative linear function with a regression equation, Y1 =  $2568.9 - 294 \, \text{Xi}$  (P <01) and a correlation coefficient,  $r = -0.86 \, (r2 = 74\%)$ . Therefore, each fortnight delay in date of sowing after the rain became constant reduced grain yield by 294 kg ha-1.

Reduction in yield associated with delayed sowing could be explained partly by the fact that some of the late sown crops did not produce flowers while others that flowered, did not set seeds. The flowering period came at the peak of the rain which might have washed off some of the pollen, thus preventing maximum seed development. This is supported Balsasubramanian (1959) who noted that heavy rainfall does interfere with




Fig 3: The effect of sowing date on grain yields of millet for 1976 to 1978




Fig.4 The quadratic effect of sowing date on grain yield of millet

seed setting. The early and late sown plants were characterized by small heads, fewer number of grain per head and fewer number of plants established (Table 1). The implication of plant establishment in yield reduction is obvious when it is considered that there is positive correlation between total yield and number of plants established (Table 5). It could also be inferred that since there was a positive correlation between grain yield and time to 50% heading, delayed heading reduced the number of days for grain filling. Grain yield in millet has been found to be related to duration and rate of filling (Egbarevba, 1979b).

# Conclusion and Recommendation

It appears from this study that growth and development of millet could be adversely affected when sown at less than 50 mm of rainfall. Sowing too late into the rainy season also has adverse effect on growth because of the resultant decrease in heat unit. Deviation from the optimum sowing time as determined by when rainfall becomes constant, will lead to a decrease in grain yield. The longer the delay, the greater the reduction in yield. The time the rain becomes constant is generally dependent on its onset and thus, the May sowing date found to be most suitable in this study, may not always be the ideal. The experience of 1977 in the present study is a case in point. A rule of thumb, then, is that millet should be sown after the first 5 — 8 rain days of the season provided the spread is not more than three weeks. This will help eliminate some of the failures encountered by farmers when they sow with the first rain.

# Acknowledgement

This work forms part of the approved research programme of the Institute for Agricultural Research, Samaru. The Director's permission to publish it is gratefully acknowledged. Mr. S. Ukata's assistance in collecting the date is also greatly acknowledged.

## References

- Amold, C. Y. (1960. Maximum-minimum temperatures as a basis for computing heat units. *Proc. Am. Soc. hort. Sci.* 76: 682-692.
- Balasubramanian, C. 1959. Crop weather relationship in rainfed sorghum. *Indian J. Agron.* 3: 247 251.
- Bonaparte, E.E.N.A. and Brawn, R. I. 1976. Effects of plant density and planting date on leaf number and some developmental events in Corn. *Can. J. Plant Sci.* 56: 691 698.
- Bunting, E. S. 1968. The influence of date of sowing on development and yield of maize in England. *J. Agric. Sci. Cambridge*,71: 117–125.
- Egharevba, P. N.: 1979a. How farmers can improve their millet yield. *Noma* (2(2): 3-6).
- Egharevba, P. N. (in press). Grain filling period as it affects yield in millet. Cereal Res. Comm.
- Evans, W. F.: and Stickler, F. C. 1961. Grain Sorghum seed germination under moisture and temperature stress. *Agron. J.* 53: 369 372.
- Genter, C. F. and Jones. G. D. 1970. Planting date and growing season effects and interactions on growth and yield of maize. *Agron. J.* 62: 760–761.
- Jones, M. J. 1973. Time of application of nitrogen fertilizer to maize at Samaru, Nigeria. *Expl. Agric. 9*: 113 120.
- Kassam, A. H. and Andrews D. J. 1975. Effects of sowing date on growth, development and yield of photosensitive sorghum at Samaru, northern Nigeria, *Expl. Agric.* 11: 227 240.
- Millington, A. J. Whiting, M.I.K, Williams, W.T. and Boundym C.A.P. 1977. The effects of sowing date on the growth and yield of three sorghum cultivars in the Ord River Valley. 1 Agronomic aspects. Aust. J. Agric. Res.; 28: 269 379.
- Stickler, F. C. and Pauli, A.W. 1961. Influence of date of planting on yield and yield components in grain sorghum. *Agron. J.* 53: 20–22.
- Zuber, M. S. :1966. Date of planting studies with Corn. Mo. Agric. Exp. Sta. Bull., 818.
- Zuber, M. S. 1968. Date of planting studies with Corn in Central Missouri. *Md. Agric. Expt. Sta. Bull.* 832.